

Übung: Molecular Dynamics Simulation of a Protein

SS 2012 Übung zu Struktur und Simulation

14-May-2012

1. Timeplan	1
2. Introduction	2
2.1. Calculations.....	2
3. Choosing your protein and temperature	3
4. Gromacs calculations	3
4.1. First MD run.....	4
4.2. Second MD run	5
5. Simple analysis	6
5.1. Energies.....	6
5.2. Fluctuations	6
6. Exchange of information.....	7
7. Assignment.....	7
8. Background and reference notes	10
8.1. Gromacs and the commands	10
8.2. The batch system.....	10
9. chimera / displaying structures / movies	11

1. Timeplan

* May 15: set up simulations and put them in the batch system

* May 22: analysis of

- * energy

- * size

- * correlation function

* Jun 5: exchange of results between different people

A brief report should be handed in by 12 Jun.

2. Introduction

It is assumed that

- * you remember how to get a file of coordinates from the protein databank (www.rcsb.org).
- * you can produce a simple x - y plot
- * you remember the program chimera, but see the notes on page 11.

The aim is to have different people run simulations at different temperatures, to see when a protein falls apart. We work on two different proteins and they will behave differently. This means we need the results from different people to write the report.

There is much room for making typing mistakes and everybody makes typing mistakes. The various programs will print an error message if something did not work. Please check the output carefully. If you see very strange results, it almost certainly means you have typed something wrong. Some people will see strange results.

2.1. Calculations

The simulations are rather realistic, but we will make a drastic simplification so as to save computer time. We will run on proteins *in vacuo*. There will be no explicit solvent. This saves more than an order of magnitude CPU time. The model uses united atoms (hydrogens merged into their carbons), but polar hydrogens are treated explicitly. There are lots of parameters and unfamiliar terms in the documentation. Either ignore them or ask what they mean.

The MD calculations will use a time step of 0.002 ps (2×10^{-15} s). You have to know this when labelling the x -axis on the plots. Each 500 steps = 1 ps.

The steps will be

- * convert files to gromacs format
- * run an energy minimisation
- * run a very short MD simulation for equilibration and not analyse anything
- * run a longer simulation in the batch queue and go home
- * do some analysis of the runs and exchange results with people who worked at different temperatures

3. Choosing your protein and temperature

To choose which protein you will simulate, add together the first letters of your first name (TOM = 20 + 15 + 13 = 48). If the number is even, use protein 2a3d, else use protein 1omb. To get the temperature you should simulate at, add together the letters of your last name like above and take that number modulo 7. If your number is n, then choose the n-th (start to count at 0) temperature from this list: 200 K, 250 K, 300 K, 350 K, 400 K, 500 K, 600 K.

4. Gromacs calculations

You are responsible for giving your input and output files sensible names. The ones given here are suggestions. If you are working on protein 1omb, you would want to replace each 2a3d below with 1omb.

Start by finding the coordinates for your protein and save them in a filename like 2a3d.pdb or 1omb.pdb.

First one removes unwanted hydrogens and adds polar hydrogens if necessary. If your protein is 2a3d, then :

```
~torda/gromacs/bin/pdb2gmh \
    -ignh -ff G43b1 -f 2a3d.pdb -o 2a3d_start.pdb -p 2a3d
```

This will put starting coordinates in 2a3d_start.pdb. The "G43b1" tells gromacs to use a united-atom force field with parameters suited to *in vacuo* simulations. 2a3d.top contains description (topology) of the molecule.

Next, one has to make input files for the simulation or minimization programs. For this, one uses "grompp". This is a pre-processor which checks if the input is sensible, writes a description of the molecule and then a long input file for the next step.

Copy the files

~torda/uebung_gromacs/*.mdp to your working directory. Open em.mdp in an editor and

* set nsteps = 50000 (the maximum number of steps of steepest descent minimisation)

* set emtol = 20 (set the convergence criterion)

Then

```
~torda/gromacs/bin/grompp \
```

```
-f em.mdp -c 2a3d_start.pdb \
-p 2a3d.top -po 2a3d_em -o 2a3d_em
```

The switches tell gromacs to run an energy minimising calculation using the steepest descents method for a maximum of 50000 steps.

Now you should have 2a3d_em.tpr and 2a3d_em.mdp which contain enough information for an energy minimisation calculation. Finally run the energy minimisation calculation:

```
~torda/gromacs/bin/mdrun \
-s 2a3d_em -o 2a3d_em -c 2a3d_em.pdb -g 2a3d_em -e 2a3d_em
```

What should happen is that the coordinates from 2a3d_start.pdb will be read, minimised and the new, lower energy coordinates written to 2a3d_em.pdb. The information about energies should be in the log file, 2a3d_em.log.

Read the log file and note

```
* how many steps were
run
* the initial energy      } look at potential energy, there is no kinetic energy in an energy
* the final energy       } minimisation
```

You might also want to use chimera to compare the starting and finishing coordinates. They are probably rather similar.

4.1. First MD run

The energy minimisation calculation will have removed some high energy problems such as bad contacts (atoms too close to each other). Now, the molecule can be relaxed further by a short MD run. We will not be doing much analysis of this step. The aim is to relax the coordinates.

If you copied the files *.mdp as described above, you should find a file called md1.mdp. Edit this file. Set the reference temperature `ref_t` and initial temperature `gen_temp` to 250 K. Set the number of steps `nsteps` to 10 000. As before, we need two commands to run the simulation.

```
~torda/gromacs/bin/grompp \
-c 2a3d_em.pdb -f md1.mdp -p 2a3d -po 2a3d_md1 -o 2a3d_md1
```

then

```
~torda/gromacs/bin/mdrun \
  -s 2a3d_md1 -o 2a3d_md1 -e 2a3d_md1 -c 2a3d_md1.pdb -g
2a3d_md1
```

which should produce

```
* 2a3d_md1.edr (energies)
* 2a3d_md1.log (log file to be read)
* 2a3d_md1.pdb (final coordinates)
* 2a3d_md1.trr (trajectory)
```

Now, you have to stop and collect some information necessary for the report.

- start chimera and look at the initial PDB file, the coordinates after energy minimising and the coordinates after the short MD run. It should be easy to superimpose the files and see if the structure has changed much at each stage. Your trajectory is very short, but you may also like to have a look at the trajectory reading features in chimera.
- from the log file, note down the initial and final energies (total, potential and kinetic) as well as the average and the root mean square (rms) fluctuations

4.2. Second MD run

The system (protein) may not be truly equilibrated, but it should roughly fit into the force field. Now, we run a second, longer simulation. The inputs will be different for different people. We will run the two steps grompp and mdrun, but the second should be run as a batch job. See the section on page 10. Edit md2.mdp. Set the temperature `ref_t` to the temperature that was given to you on page **Error! Bookmark not defined.** Then edit md2.mdp and set `nsteps` to a very small number like 20.

```
~torda/gromacs/bin/grompp \
  -c 2a3d_md1 -f md2.mdp -p 2a3d -po 2a3d_md2 -o 2a3d_md2
~torda/gromacs/bin/mdrun \
  -s 2a3d_md2 -o 2a3d_md2 -c 2a3d_md2.pdb -e 2a3d_md2 \
  -g 2a3d_md2
```

Check that the commands have worked. For example, do you have a `2a3d_md2.tpr` and the corresponding log file ? If so, create a script file to run the commands. Submit it to the batch system and make sure it works. If the gromacs commands are correct and you can run a command

in the batch system, edit `md2.mdp` and set `nsteps` to 250 000. Submit this long job to the batch system. Make sure the command is really running and go home.

5. Simple analysis

5.1. Energies

From your calculations, you should have files like

`xxx_md2.trr` (trajectory)

`xxx_md2.edr` (energies)

`xxxx_md2.log` (log file)

where `xxx` will depend on your protein and the names you used.

Read the log file. Note down the start, finish, average and root mean square fluctuations of the total, potential and kinetic energies. In chimera, try to look at the trajectory file.

First look at the energies from the calculations. There is a program which will write these in a format for `xmgrace`. Try

```
~torda/gromacs/bin/g_energy -f xxx.edr -o xxx.xvg
```

then

```
xmgrace xxx.xvg
```

You do not have to use `xmgrace`, but it should be convenient. The program `g_energy` will ask which energies you want. Try

Potential

Kinetic-En.

Total-Energy

Giving each answer on its own line.

5.2. Fluctuations

When looking at the properties of a trajectory, the gromacs programs will often ask which atoms you want to look at. In each case, we will choose the backbone atoms only.

It is not impossible that some people remember the formula for root mean square differences of coordinates given last semester. In an MD simulation, it is useful to compare each snapshot with the starting coordinates, to see how far the structure has moved. The formula is still

$rmsd = \left(\frac{1}{n} \sum_{i=1}^n |\vec{r}_{1,i} - \vec{r}_{2,i}|^2 \right)^{\frac{1}{2}}$ where $\vec{r}_{1,i}$ is the starting configuration and $\vec{r}_{2,i}$ a snapshot from a

trajectory. To see this from a gromacs trajectory,

```
~torda/gromacs/bin/g_rms -f xxx.trr -s xxx.tpr  
-o xxx_rms.xvg
```

Your results will not be the same as anybody else's. The "xxx" will refer to your protein and your temperature.

Next we should look at the radius of gyration of the protein. This is a measure of the size of a protein. First one calculates the centre of mass and then the root mean square difference of each atom to this point. The gromacs command is

```
~torda/gromacs/bin/g_gyrate -f xxx.trr -s xxx.tpr  
-o xxx_gyr.xvg
```

and the file, xxx_gyr.xvg can be viewed and printed with xmgrace.

Now, for the last part of the Übung, please make sure you have a plot on paper of the radius of gyration and rmsd for your protein.

6. Exchange of information

In the third week, we will attempt to exchange the numbers necessary to fill out the table on page 8. This we will do on the blackboard so we can compare and discuss the numbers and find mistakes now.

For this third week, bring a plot on paper of the radius of gyration and rmsd of your protein as a function of time. Label the plot with the name of your protein and the temperature. We will then put these through a photocopier so everybody can have a copy.

7. Assignment

Answers should be on pieces of paper, not be electronic.

All graphs must have labels on the axes with units. Any numbers in tables should have units.

Please do not quote more significant digits than is sensible. If you have a quantity such as

2355.55 ± 323 kJ mol⁻¹ (rms fluctuation), the best way to quote the energy is 2.4 × 10³ kJ mol⁻¹.

1. Write the most important details of your protein:

- * its name
- * number of residues
- * how many disulfide bridges does it have
- * very crude description of structure (mostly α -helical, mixed α/β , much irregular structure..)
- * how the structure was solved (X-ray, NMR)
- * write the temperature you did your simulation at

2.

* Build a table for each protein like:

				Temperature (K)		
	initial	after EM	MD1	200	250	...
	$\text{kJ mol}^{-1} \pm \text{rms fluctuations}$					
E_{tot}						
E_{pot}						
E_{kin}	X	X				

E_{tot} , E_{pot} , E_{kin} are the total, potential and kinetic energies respectively. There is no kinetic energy from an energy minimisation. Rms fluctuation refers to the root mean square fluctuations. All these quantities are in the xxx.log files from the calculations. For all MD runs, quote the average energy values.

* For your protein and your temperature, produce an x - y plot from the second MD simulation of total, potential and kinetic energies as a function of time. Label all axes and put in units.

During this longer simulation,

* Is there a trend in total energy ?

* Does the system seem to be equilibrated ?

* Are the fluctuations of potential and kinetic energy correlated ?

3. In chimera, compare the structure

(a) from the PDB

(b) after energy minimising

(c) after the first MD run

(d) after the second MD run

* Does the structure change much in any single step ? The answer to this question will be different for every person in the class.

* If you view the trajectory in chimera, is there any trend ? Does the structure open or close ? Do the secondary structure units remain intact ?

* Does the plot of radius of gyration or root mean square displacement (rmsd) support what you say ?

4. In some of the simulations, there will be trend in the total energy. It will have decreased. This seems to violate energy conservation. Where has the energy gone to ? This question requires the lecture material on temperature baths.

5. I make the claim that for a simulation at 300 K, a protein simulation will tend to move the protein towards a lower energy region of conformations. If you look at the average energies from the trajectories, there is no evidence for this. What would you do to try to confirm my claim ? (that the protein has moved closer to low energy regions of space).

6. Imagine that after simulating at 300 K, the average coordinates have definitely moved away from the starting conformation, but the energy does not seem to have increased. A chemist would

say that the native conformation from the protein data bank should be the lowest energy. What is a plausible explanation ?

8. Background and reference notes

8.1. Gromacs and the commands

Gromacs is one of the popular MD codes used for research projects. It has been installed under `~torda/gromacs`. There is no single program called "gromacs". Instead, there is a collection of programs in `~torda/gromacs/bin`.

Gromacs is free and can be found at www.gromacs.org. This web page also has the full documentation. We have the text for the man pages sitting in the distribution directory. Try a command like

```
cd ~torda/gromacs/man/man1
man ./grompp.1
man ./mdrun.1
```

We will also be using the commands `g_energy` and `g_rms`.

8.2. The batch system

Usually you run a job and wait for the answer. If your job will take more than a few minutes, you may not want to sit and wait. It is often more convenient to use the "batch system" on the machines. It is also more sociable than simply running in the background. Using the batch system ensures that only one job at a time will run in the background and other users can easily see which machines are busy.

The principle is :

- * you put your commands in a script file (usually `/bin/sh` or `/bin/tcsh`).
- * you submit (qsub) your script
- * the batch system will find a machine which is not too busy and run the job in the background

The batch system relies on some environment variables, so you must set some symbols. Do this by sourcing a file

```
source /opt/nlge6/zbh/common/settings.sh    for bash users
```

Source /opt/nlge6/zbh/common/settings.csh for csh or tcsh users

There are man pages for all commands such as `man qsub` and `man qstat`. Before submitting an MD job, make sure a toy example works. Create a file like `simple.sh` with some simple commands:

```
#!/bin/sh
uname -a
date
sleep 20
```

Submit the job

```
qsub -clear -q stud.q simple.sh
```

if you would like mail when the job finishes, try

```
qsub -clear -q stud.q -m e -M yourname@somehost.de simple.sh
```

Type `qstat` to see which jobs are running. When running your longer MD job, do make sure that your job is really running and has not crashed.

To run your MD job in a batch queue, put your commands in a file and submit that file instead of `simple.sh` used above.

9. chimera / displaying structures / movies

Everyone has used "chimera" last semester. It can display trajectories from simulations from a program like gromos. It is installed as `/usr/local/zbh/chimera/bin/chimera`.

If you have a trajectory of a molecule like 1omp, you will need the trajectory file `xxx.trr` and the atomic description file, `xxx.tpr`. Look for the menu entry, "tools", then "MD/Ensemble analysis". Specify the format as gromacs.